arXiv:1610.03799v2 [math.NT] 19 Oct 2016

# Some supercongruences for truncated generalized hypergeometric series ${}_3F_2$

Ji-Cai Liu

Department of Mathematics, East China Normal University,
500 Dongchuan Road, Shanghai 200241, People's Republic of China
jc2051@163.com

**Abstract.** Rodriguez-Villegas conjectured 22 supercongruences for hypergeometric Calabi-Yau manifolds of dimension $D \leq 3$. For manifolds of dimension $D = 2$, he posed four conjectural supercongruences on truncated generalized hypergeometric series ${}_3F_2$. These four conjectures were gradually confirmed by several authors. Z.-H. Sun has extended these four supercongruences in a unified form. In this paper, we aim to generalize the results of Z.-H. Sun by using some combinatorial identities involving harmonic numbers.

*Keywords*: supercongruences; generalized hypergeometric series; harmonic numbers

*MR Subject Classifications*: Primary 11A07; Secondary 33C05

## 1 Introduction

Rodriguez-Villegas [9] observed the remarkable relationship between the number of points over $F_p$ on certain Calabi-Yau manifolds and truncated generalized hypergeometric series. He conjectured numerically 22 supercongruences for hypergeometric Calabi-Yau manifolds, which can be categorized by the dimension of the manifolds.

We first define the truncated generalized hypergeometric series. For complex numbers $a_i, b_j$ and $z$, with none of the $b_j$ being negative integers or zero, the truncated generalized hypergeometric series are given by

$$ {}_rF_s\left[\begin{matrix} a_1, a_2, \cdots, a_r \\ b_1, b_2, \cdots, b_s \end{matrix}; z\right]_n = \sum_{k=0}^{n-1} \frac{(a_1)_k (a_2)_k \cdots (a_r)_k}{(b_1)_k (b_2)_k \cdots (b_s)_k} \cdot \frac{z^k}{k!}, $$

where $(a)_0 = 1$ and $(a)_k = a(a+1)\cdots(a+k-1)$ for $k \geq 1$.

For manifolds of dimension $D = 1$, Rodriguez-Villegas [9] posed 4 conjectural supercongruences, which were first confirmed by Mortenson [6, 7]. For dimension $D = 2$, another 4 conjectural supercongruences can be written in the form

$$ {}_3F_2\left[\begin{matrix} \frac{1}{2}, -a, a+1 \\ 1, 1 \end{matrix}; 1\right]_p \equiv c(p) \pmod{p^2}, $$

where $a=-\frac{1}{2},-\frac{1}{3},-\frac{1}{4},-\frac{1}{6}$, $p$ is an odd prime and $c(p)$ is the $p$-th Fourier coefficient of a weight three modular form on a congruence subgroup of $SL(2,\mathbb{Z})$. In particular,

$$ {}_3F_2\left[\begin{matrix}\frac{1}{2},\frac{1}{2},\frac{1}{2}\\ 1,1\end{matrix};1\right]_p\equiv 0 \pmod{p^2} \quad \text{for } p\equiv 3 \pmod 4, \tag{1.1}$$

$$ {}_3F_2\left[\begin{matrix}\frac{1}{2},\frac{1}{3},\frac{2}{3}\\ 1,1\end{matrix};1\right]_p\equiv 0 \pmod{p^2} \quad \text{for } p\equiv 5 \pmod 6, \tag{1.2}$$

$$ {}_3F_2\left[\begin{matrix}\frac{1}{2},\frac{1}{4},\frac{3}{4}\\ 1,1\end{matrix};1\right]_p\equiv 0 \pmod{p^2} \quad \text{for } p\equiv 5,7 \pmod 8, \tag{1.3}$$

$$ {}_3F_2\left[\begin{matrix}\frac{1}{2},\frac{1}{6},\frac{5}{6}\\ 1,1\end{matrix};1\right]_p\equiv 0 \pmod{p^2} \quad \text{for } p\equiv 3 \pmod 4. \tag{1.4}$$

It should be mentioned that (1.1) was originally conjectured by Beukers and Stienstra [3] and first proved by van Hamme [16]. Supercongruences (1.2)-(1.4) were finally confirmed by Z.-W. Sun [15].

By studying the generalized Legendre polynomials, Z.-H. Sun [14, Theorem 2.5] have extended (1.1)-(1.4) in a unified form as follows.

**Theorem 1.1** *(Z.-H. Sun) Let $p$ be an odd prime. For any $p$-adic integer $a$ with $\langle a\rangle_p\equiv 1 \pmod 2$, we have*

$$ {}_3F_2\left[\begin{matrix}\frac{1}{2},-a,a+1\\ 1,1\end{matrix};1\right]_p\equiv 0 \pmod{p^2}, \tag{1.5}$$

*where $\langle a\rangle_p$ denotes the least non-negative integer $r$ with $a\equiv r \pmod p$.*

Letting $a=-\frac{1}{2},-\frac{1}{3},-\frac{1}{4},-\frac{1}{6}$ in (1.5) reduce to (1.1)-(1.4).

In this paper, we aim to generalize (1.5) by using an essentially different method. Our proof is based on some combinatorial identities involving harmonic numbers.

**Theorem 1.2** *Suppose $p$ is an odd prime and $n$ is a positive integer. For any $p$-adic integer $a$ with $\langle a\rangle_p\equiv 1 \pmod 2$, we have*

$$ {}_3F_2\left[\begin{matrix}\frac{1}{2},-a,a+1\\ 1,1\end{matrix};1\right]_{np}\equiv 0 \pmod{p^2}. \tag{1.6}$$

In the next section, we first prove some important lemmas. The proof of Theorem 1.2 will be given in Section 3.

## 2 Some lemmas

**Lemma 2.1** *Let $H_n=\sum_{k=1}^{n}\frac{1}{k}$ be the $n$-th harmonic number. For any odd positive integer $n$, we have*

$$\sum_{k=0}^{n}\binom{2k}{k}^2\binom{n+k}{2k}\left(-\frac{1}{4}\right)^k=0, \tag{2.1}$$

$$\sum_{k=0}^{n}\binom{2k}{k}^2\binom{n+k}{2k}\left(-\frac{1}{4}\right)^k(2H_{2k}-3H_k)=0, \tag{2.2}$$

$$\sum_{k=0}^{n}\binom{2k}{k}^2\binom{n+k}{2k}\left(-\frac{1}{4}\right)^k(2H_{n+k}-H_k)=0. \tag{2.3}$$

*Proof.* Note the following identity [2, (1), pp. 16]:

$$\begin{aligned}&{}_3F_2\left[\begin{matrix}a,b,c\\ \frac{1}{2}(a+b+1),2c\end{matrix};1\right]\\ &=\frac{\Gamma\left(\frac{1}{2}\right)\Gamma\left(\frac{1}{2}+c\right)\Gamma\left(\frac{1}{2}+\frac{1}{2}a+\frac{1}{2}b\right)\Gamma\left(\frac{1}{2}-\frac{1}{2}a-\frac{1}{2}b+c\right)}{\Gamma\left(\frac{1}{2}+\frac{1}{2}a\right)\Gamma\left(\frac{1}{2}+\frac{1}{2}b\right)\Gamma\left(\frac{1}{2}-\frac{1}{2}a+c\right)\Gamma\left(\frac{1}{2}-\frac{1}{2}b+c\right)},\end{aligned} \tag{2.4}$$

where the gamma function [1, (1.1.5), pp. 3] is given by $\Gamma(x)=\lim_{k\to\infty}k!k^{x-1}/(x)_k$.

Letting $a=-n, b=n+1$ and $c=\frac{1}{2}$ in (2.4) reduces to

$${}_3F_2\left[\begin{matrix}-n,n+1,\frac{1}{2}\\ 1,1\end{matrix};1\right]=\frac{\Gamma\left(\frac{1}{2}\right)^2}{\Gamma\left(\frac{1-n}{2}\right)^2\Gamma\left(\frac{n+2}{2}\right)^2}.$$

Since $(1-n)/2$ is always a non-positive integer, we have $1/\Gamma\left(\frac{1-n}{2}\right)^2=0$, and so

$${}_3F_2\left[\begin{matrix}-n,n+1,\frac{1}{2}\\ 1,1\end{matrix};1\right]=0,$$

for any odd positive integer $n$. Noting that

$$\frac{\left(\frac{1}{2}\right)_k}{(1)_k}=\frac{\binom{2k}{k}}{4^k}, \tag{2.5}$$

$$\frac{(-n)_k(n+1)_k}{(1)_k^2}=(-1)^k\binom{2k}{k}\binom{n+k}{2k}, \tag{2.6}$$

we are led to (2.1).

Let $A_n, B_n$ and $C_n$ denote the numbers

$$A_n=\sum_{k=0}^{2n-1}\binom{2k}{k}^2\binom{2n-1+k}{2k}\left(-\frac{1}{4}\right)^k H_k,$$

$$B_n=\sum_{k=0}^{2n-1}\binom{2k}{k}^2\binom{2n-1+k}{2k}\left(-\frac{1}{4}\right)^k H_{2k},$$

and

$$C_n=\sum_{k=0}^{2n-1}\binom{2k}{k}^2\binom{2n-1+k}{2k}\left(-\frac{1}{4}\right)^k H_{2n-1+k},$$

respectively. Using the software package developed by C. Schneider [10], we find that $A_n, B_n$ and $C_n$ satisfy the same recurrence:

$$\begin{aligned}&4n^2(2n+1)^2(4n+5)x_n-(4n+3)(32n^4+96n^3+104n^2+48n+7)x_{n+1}\\&+4(n+1)^2(2n+3)^2(4n+1)x_{n+2}=0.\end{aligned}$$

It is easy to verify that $3A_n=2B_n=6C_n$ for $n=1,2$. This implies that $3A_n=2B_n=6C_n$ for any positive integer $n$. □

**Lemma 2.2** *Suppose $p$ is an odd prime. For any $p$-adic integer $a$ with $\langle a\rangle_p\equiv 1 \pmod 2$, we have*

$$\begin{aligned}&\sum_{k=0}^{p-1}\binom{2k}{k}^2\binom{a+k}{2k}\left(-\frac{1}{4}\right)^k\sum_{i=1}^{k}\frac{1}{-a-1+i}\\&\equiv\sum_{k=0}^{p-1}\binom{2k}{k}^2\binom{a+k}{2k}\left(-\frac{1}{4}\right)^k\sum_{i=1}^{k}\frac{1}{a+i}\\&\equiv\frac{1}{2}\sum_{k=0}^{p-1}\binom{2k}{k}^2\binom{a+k}{2k}\left(-\frac{1}{4}\right)^k H_k\pmod p.\end{aligned}\tag{2.7}$$

*Proof.* From (2.1) and (2.3) we have

$$\sum_{k=0}^{n}\binom{2k}{k}^2\binom{n+k}{2k}\left(-\frac{1}{4}\right)^k(2H_{n+k}-2H_n-H_k)=0,$$

for any odd positive integer $n$, which is equivalent to

$$\sum_{k=0}^{n}\binom{2k}{k}^2\binom{n+k}{2k}\left(-\frac{1}{4}\right)^k\left(\sum_{i=1}^{k}\frac{1}{n+i}-\frac{1}{2}H_k\right)=0.\tag{2.8}$$

Since $\langle a\rangle_p$ is an odd positive integer, it follows that

$$\begin{aligned}&\sum_{k=0}^{p-1}\binom{2k}{k}^2\binom{a+k}{2k}\left(-\frac{1}{4}\right)^k\sum_{i=1}^{k}\frac{1}{a+i}\\&\equiv\sum_{k=0}^{\langle a\rangle_p}\binom{2k}{k}^2\binom{\langle a\rangle_p+k}{2k}\left(-\frac{1}{4}\right)^k\sum_{i=1}^{k}\frac{1}{\langle a\rangle_p+i}\\&=\frac{1}{2}\sum_{k=0}^{\langle a\rangle_p}\binom{2k}{k}^2\binom{\langle a\rangle_p+k}{2k}\left(-\frac{1}{4}\right)^k H_k\quad\text{(by (2.8))}\\&\equiv\frac{1}{2}\sum_{k=0}^{p-1}\binom{2k}{k}^2\binom{a+k}{2k}\left(-\frac{1}{4}\right)^k H_k\pmod p.\end{aligned}$$

Let $b$ denote the positive integer $b = p - \langle a \rangle_p$. It is clear that $a \equiv -b \pmod{p}$ and $b-1$ is an odd positive integer. It follows that

$$\begin{aligned}&\sum_{k=0}^{p-1}\binom{2k}{k}^2\binom{a+k}{2k}\left(-\frac{1}{4}\right)^k\sum_{i=1}^{k}\frac{1}{-a-1+i}\\&\equiv\sum_{k=0}^{p-1}\binom{2k}{k}^2\binom{-b+k}{2k}\left(-\frac{1}{4}\right)^k\sum_{i=1}^{k}\frac{1}{b-1+i}\pmod{p}.\end{aligned}\tag{2.9}$$

Since $\binom{-b+k}{2k} = \binom{b-1+k}{2k}$, by (2.8) and (2.9) we get

$$\begin{aligned}\text{LHS (2.9)} &\equiv \sum_{k=0}^{b-1}\binom{2k}{k}^2\binom{b-1+k}{2k}\left(-\frac{1}{4}\right)^k\sum_{i=1}^{k}\frac{1}{b-1+i}\\&= \frac{1}{2}\sum_{k=0}^{b-1}\binom{2k}{k}^2\binom{b-1+k}{2k}\left(-\frac{1}{4}\right)^k H_k \quad \text{(by (2.8))}\\&\equiv \frac{1}{2}\sum_{k=0}^{p-1}\binom{2k}{k}^2\binom{-a-1+k}{2k}\left(-\frac{1}{4}\right)^k H_k\\&= \frac{1}{2}\sum_{k=0}^{p-1}\binom{2k}{k}^2\binom{a+k}{2k}\left(-\frac{1}{4}\right)^k H_k \pmod{p}.\end{aligned}$$

This completes the proof of (2.7). □

# 3 Proof of Theorem 1.2

Noting (2.5) and (2.6), we can rewrite (1.6) as

$$\sum_{k=0}^{np-1}\binom{2k}{k}^2\binom{a+k}{2k}\left(-\frac{1}{4}\right)^k \equiv 0 \pmod{p^2}.$$

We first prove that for any non-negative integer $r$

$$\sum_{k=rp}^{(r+1)p-1}\binom{2k}{k}^2\binom{a+k}{2k}\left(-\frac{1}{4}\right)^k \equiv 0 \pmod{p^2},\tag{3.1}$$

which is equivalent to

$$\sum_{k=0}^{p-1}\binom{2k+2rp}{k+rp}^2\binom{a+k+rp}{2k+2rp}\left(-\frac{1}{4}\right)^{k+rp} \equiv 0 \pmod{p^2}.\tag{3.2}$$

Applying the the following congruence [4, (2.1)]:

$$\binom{2rp+2k}{rp+k} \equiv \binom{2r}{r}\binom{2k}{k}(1+2rp(H_{2k}-H_k)) \pmod{p^2},$$

we immediately get

$$\binom{2rp+2k}{rp+k}^2 \equiv \binom{2r}{r}^2\binom{2k}{k}^2(1+4rp(H_{2k}-H_k)) \pmod{p^2}. \tag{3.3}$$

By (3.3) and the fact that $\binom{2k}{k}^2 \equiv 0 \pmod{p^2}$ for $\frac{p-1}{2} < k \le p-1$, we have

$$\begin{aligned}&\text{LHS (3.2)}\\ &\equiv \sum_{k=0}^{\frac{p-1}{2}} \binom{2r}{r}^2\binom{2k}{k}^2\binom{a+k+rp}{2k+2rp}\left(-\frac{1}{4}\right)^{k+rp}(1+4rp(H_{2k}-H_k)) \pmod{p^2}. \end{aligned}\tag{3.4}$$

Let $\delta$ denote the number $\delta = (a-\langle a\rangle_p)/p$. It is clear that $\delta$ is a $p$-adic integer and $a = \langle a\rangle_p + \delta p$. Note that

$$\begin{aligned}&\binom{a+k+rp}{2k+2rp}\\ &= \binom{\langle a\rangle_p + k + (\delta+r)p}{2k+2rp}\\ &= \binom{\langle a\rangle_p + (\delta+r)p}{2rp}\prod_{i=1}^{k}(\langle a\rangle_p + (\delta+r)p+i)\prod_{i=1}^{k}(\langle a\rangle_p + (\delta-r)p+1-i)\prod_{i=1}^{2k}(2rp+i)^{-1}.\end{aligned}$$

Since

$$(2rp+i)^{-1} \equiv \frac{1}{i} - \frac{2rp}{i^2} \pmod{p^2} \quad \text{for } 1 \le i \le p-1,$$

it follows that for $0 \le k \le \frac{p-1}{2}$

$$\begin{aligned}\binom{a+k+rp}{2k+2rp} &\equiv \binom{\langle a\rangle_p + (\delta+r)p}{2rp}\\ &\quad\times \prod_{i=1}^{k}(\langle a\rangle_p + (\delta+r)p+i)\prod_{i=1}^{k}(\langle a\rangle_p + (\delta-r)p+1-i)\prod_{i=1}^{2k}(\frac{1}{i}-\frac{2rp}{i^2})\\ &\equiv \binom{\langle a\rangle_p + (\delta+r)p}{2rp}\binom{\langle a\rangle_p + k}{2k}\\ &\quad\times\left(1+p\left(\sum_{i=1}^{k}\frac{r+\delta}{\langle a\rangle_p + i} + \sum_{i=1}^{k}\frac{r-\delta}{-\langle a\rangle_p - 1 + i} - 2rH_{2k}\right)\right) \pmod{p^2}.\end{aligned}\tag{3.5}$$

Combining (3.4) and (3.5), we obtain

$$\begin{aligned}\text{LHS (3.2)} \equiv &\left(-\frac{1}{4}\right)^{rp}\binom{2r}{r}^2\binom{\langle a\rangle_p+(\delta+r)p}{2rp}\sum_{k=0}^{\frac{p-1}{2}}\binom{2k}{k}^2\binom{\langle a\rangle_p+k}{2k}\left(-\frac{1}{4}\right)^k\\ &\times\left(1+p\left(\sum_{i=1}^{k}\frac{r+\delta}{\langle a\rangle_p+i}+\sum_{i=1}^{k}\frac{r-\delta}{-\langle a\rangle_p-1+i}+2rH_{2k}-4rH_k\right)\right)\pmod{p^2}.\end{aligned}\tag{3.6}$$

Note that $\binom{2k}{k}^2\equiv 0 \pmod{p^2}$ for $\frac{p-1}{2}<k\le p-1$ and $\langle a\rangle_p$ is an odd positive integer. It follows from (2.1) that

$$\begin{aligned}&\sum_{k=0}^{\frac{p-1}{2}}\binom{2k}{k}^2\binom{\langle a\rangle_p+k}{2k}\left(-\frac{1}{4}\right)^k\\ &\equiv\sum_{k=0}^{\langle a\rangle_p}\binom{2k}{k}^2\binom{\langle a\rangle_p+k}{2k}\left(-\frac{1}{4}\right)^k\\ &=0\pmod{p^2}.\end{aligned}\tag{3.7}$$

On the other hand, using (2.7) and (2.2) we have

$$\begin{aligned}&\sum_{k=0}^{\frac{p-1}{2}}\binom{2k}{k}^2\binom{\langle a\rangle_p+k}{2k}\left(-\frac{1}{4}\right)^k\left(\sum_{i=1}^{k}\frac{r+\delta}{\langle a\rangle_p+i}+\sum_{i=1}^{k}\frac{r-\delta}{-\langle a\rangle_p-1+i}+2rH_{2k}-4rH_k\right)\\ &\equiv\sum_{k=0}^{\langle a\rangle_p}\binom{2k}{k}^2\binom{\langle a\rangle_p+k}{2k}\left(-\frac{1}{4}\right)^k(2rH_{2k}-3rH_k)\\ &=0\pmod{p}.\end{aligned}\tag{3.8}$$

Then the proof of (3.2) directly follows from (3.6)-(3.8).

Taking the sum over $r$ from 0 to $n-1$ on both sides of (3.1) gives

$$\sum_{k=0}^{np-1}\binom{2k}{k}^2\binom{a+k}{2k}\left(-\frac{1}{4}\right)^k\equiv 0\pmod{p^2}.$$

This completes the proof of Theorem 1.2.

# References


[1] G. E. Andrews, R. Askey and R. Roy, Special Functions, Cambridge University Press, 1999.

[2] W.N. Bailey, Generalized Hypergeometric Series, Stechert-Hafner, 1964.

[3] F. Beukers and J. Stienstra, On the Picard-Fuchs equation and the formal Brauer group of certain elliptic K3 surfaces, Math. Ann. 271 (1985), 269–304.

[4] J.-C. Liu, On two conjectural supercongruences of Apagodu and Zeilberger, J. Difference Equ. Appl., online, DOI: 10.1080/10236198.2016.1245295.

[5] D. McCarthy, On a supercongruence conjecture of Rodriguez-Villegas, Proc. Amer. Math. Soc. 140 (2012), 2241–2254.

[6] E. Mortenson, A supercongruence conjecture of Rodriguez-Villegas for a certain truncated hypergeometric function, J. Number Theory 99 (2003), 139–147.

[7] E. Mortenson, Supercongruences between truncated ${}_2F_1$ hypergeometric functions and their Gaussian analogs, Trans. Amer. Math. Soc. 355 (2003), 987–1007.

[8] E. Mortenson, Supercongruences for truncated ${}_{n+1}F_n$ hypergeometric series with applications to certain weight three newforms, Proc. Amer. Math. Soc. 133 (2005), 321–330.

[9] F. Rodriguez-Villegas, Hypergeometric families of Calabi-Yau manifolds, in: Calabi-Yau Varieties and Mirror Symmetry (Toronto, ON, 2001), Fields Inst. Commun., 38, Amer. Math. Soc., Providence, RI, 2003, 223–231.

[10] C. Schneider, An implementation of Karrs summation algorithm in mathematica, Sém. Lothar. Combin. S43b (1999), 1–10.

[11] Z.-H. Sun, Congruences involving $\binom{2k}{k}^2\binom{3k}{k}$, J. Number Theory 133 (2013), 1572–1595.

[12] Z.-H. Sun, Congruences concerning Legendre polynomials II, J. Number Theory 133 (2013), 1950–1976.

[13] Z.-H. Sun, Legendre polynomials and supercongruences, Acta Arith. 159 (2013), 169–200.

[14] Z.-H. Sun, Generalized Legendre polynomials and related supercongruences, J. Number Theory 143 (2014), 293–319.

[15] Z.-W. Sun, On sums involving products of three binomial coefficients, Acta Arith. 156 (2012), 123–141.

[16] L. van Hamme, Proof of a conjecture of Beukers on Apéry numbers, Proceedings of the conference on p-adic analysis (Houthalen, 1987), 189–195.